


Recommended Vaccine Schedule FOR Foals

Based on American Association of Equine Practitioners (AAEP) Vaccination Guidelines¹

Your foal initially gets maternal antibodies from the mare, but as they begin to wane, it's important to help protect your foal from common diseases with the appropriate vaccines. In their first year, foals require several doses of both core and risk-based vaccinations to develop protection.



	Disease	Schedule
CORE VACCINATIONS: Protection from diseases that are endemic to a region, those with potential public health significance, required by law, highly infectious, or pose a risk of severe disease and death.	Tetanus	First vaccination at 4-6 months of age
	West Nile Virus (WNV)	Second dose 4-6 weeks after the first dose
	Eastern Equine Encephalomyelitis (EEE)	Booster 10-12 months of age and annually thereafter
	Western Equine Encephalomyelitis (WEE)	
	Rabies	
RISK-BASED VACCINATIONS: Protection from diseases which your horse may be more likely to encounter depending on where you live, your travel plans, and amount of socialization selected for use based on risk assessment performed by, or in consultation with, a licensed veterinarian.	Equine Influenza (EIV)	First vaccination at 4-6 months of age
	Equine Herpesvirus (EHV-1 and EHV-4)	Second dose 3-4 weeks after first dose Booster 10-12 months of age and annually thereafter Horses at increased risk of exposure may receive booster every 6 months
	Leptospirosis	First vaccination at 6 months of age Second dose 3-4 weeks after first dose Annual revaccination
	Strangles <i>Streptococcus equi</i>	First dose at 9 months of age Second dose 3 weeks after first dose Third dose at 11-12 months of age
	Equine Arteritis Virus (EAV)	Colt (male) foals*: Single dose at 6-12 months of age

***NOTE:** Prior to initial vaccination, colt (male) foals should undergo serologic testing and be confirmed negative for antibodies to EAV. Testing should be performed shortly prior to, or preferably at, the time of vaccination. As foals can carry colostral derived antibodies to EAV for up to 6 months, testing and vaccination should **not** be performed prior to 6 months of age.¹

Consult your veterinarian to develop a vaccination program to protect your foal.

*Although FLUVAC INNOVATOR EHV-4/1 is safe for use in foals 3 months of age or older, efficacy has not been established in foals under 10 months of age.

¹ The American Association of Equine Practitioners. Vaccination Guidelines. Available at <https://aaep.org/guidelines/vaccination-guidelines>. Accessed on June 26, 2023.

All trademarks are the property of Zoetis Services LLC or a related company or a licensor unless otherwise noted. © 2023 Zoetis Services LLC. All rights reserved. COR-00230

